

[illegible]

5

10

15

20

25

30

a

35

sub 7
C2

a

5. The process according to ~~any of claims 1 to 4~~,^{claim 1} wherein said protective peptide has 30 to 200 amino acid residues.

a

5

6. The process according to ~~any of claims 1 to 5~~,^{claim 1} wherein an ion exchange resin is used in the purification process.

7. The process according to claim 6, wherein said ion exchange resin is a cation exchange resin.

a

10

8. The process according to ~~any of claims 1 to 5~~,^{claim 1} wherein a reverse phase chromatography or a hydrophobic chromatography is used in the purification process.

a

15

9. The process according to ~~any of claims 1 to 8~~,^{claim 1} wherein a surfactant and/or a salt are added in at least one of steps (1) to (5) to maintain the solubility of the peptide of interest.

c

10. The process according to ~~any of claims 1 to 9~~,^{claim 1} wherein the host cell is a prokaryotic cell or a eukaryotic cell.

20

11. The process according to claim 10, wherein the host cell is Escherichia coli.

a

12. The process according to ~~any of claims 1 to 11~~,^{claim 1} wherein the isoelectric point of the peptide of interest having a helper peptide added thereto is 8 to 12.

a

25

13. The process according to ~~any of claims 1 to 11~~,^{claim 1} wherein the peptide of interest is an amidated peptide.

a

14. The process according to ~~any of claims 1 to 11~~,^{claim 1} wherein the peptide of interest is a GLP-1 derivative having an insulinotropic activity.

30

15. The process according to claim 14, wherein the GLP-1 derivative having an insulinotropic activity that has a helper peptide added thereto has an isoelectric point of 8 to 12.

a

35

16. The process according to claim 14 ~~or 15~~, wherein the GLP-1 derivative having an insulinotropic activity has an isoelectric point of 4.5 to 9.0.

a

17. The process according to claim 14 ~~or 15~~, wherein the GLP-1 derivative having an insulinotropic

activity has an isoelectric point of 5.5 to 7.5.

a
a 18. The process according to ^{claim 12} ~~any of claims 12 to 17~~, wherein an ion exchange resin is used in the purification process.

5 19. The process according to claim 18, wherein said ion exchange resin is a cation exchange resin.

a
a 20. The process according to ^{claim 12} ~~any of claims 12 to 17~~, wherein a reverse phase chromatography or a hydrophobic chromatography is used in the purification process.

10 21. The process according to ^{claim 12} ~~any of claims 12 to 17~~, wherein a surfactant and/or a salt is added to maintain the solubility of the peptide of interest.

a
15 a 22. The process according to ^{claim 14} ~~any of claims 14 to 21~~, wherein the purity of the GLP-1 derivative obtained having an insulinotropic activity is 98% or greater.

a
20 23. The process according to ^{claim 1} ~~any of claims 1 to 22~~, wherein the content of endotoxin in the final purified product is not greater than 0.03 units/mg.

a
a 24. A pharmaceutical composition for treatment of diabetes mellitus comprising as an active ingredient a GLP-1 derivative having an insulinotropic activity, obtained from the process according to ^{claim 14} ~~any of claims 14 to 23~~.

25 25. An expression vector comprising a nucleotide sequence encoding a peptide of interest that has a helper peptide added thereto, or a fusion protein that has a protective peptide further added to the peptide of interest that has a helper peptide added thereto.

30 26. A prokaryotic or a eukaryotic cell transformed with an expression vector comprising a nucleotide sequence encoding a peptide of interest that has a helper peptide added thereto, or a fusion protein that has a protective peptide further added to the peptide of interest that has a helper peptide added thereto.

35 27. The cell according to claim 26, wherein the host cell is Escherichia coli.

add
add